

Significativement plus faible que les autres interactions protéines-protéines aux conséquences biologiques



Heidi Hørig; *Expert Reviews in Molecular Medicine*, 2000

Importance de l'affinité du TCR pour l'activation lymphocytaire (1)

CD8⁺ T Cell Activation Is Governed by TCR-Peptide/MHC Affinity, Not Dissociation Rate¹

Shaomin Tian,^{*} Robert Maile,^{*1} Edward J. Collins,^{*1} and Jeffrey A. Frelinger^{2*}

CD8⁺ TCR-transgénique (P14 = TCR Va2 Vb8)
Ag = GP33-41 / LCMV
Or GP33-41 mutants (C9L, K1M/C9M)

Table I. Peptide variants of gp33

Mutation Site	Sequence	Peptide Mutant
None	SAVYNFATC	gp33
P1	SAVYNFATC	K1A, K1D, K1E, K1F, K1L, K1N, K1R, K1S, K1V, K1W, K1Y
P2	SAVYNFATC	A2D, A2E, A2F, A2G, A2K, A2L, A2N, A2R, A2S, A2T, A2V, A2W, A2Y
P3	SAVYNFATC	V3A, V3D, V3E, V3F, V3G, V3H, V3L, V3N, V3R, V3S, V3T, V3W, V3Y
P4	SAVYNFATC	Y4A, Y4C
P6	SAVYNFATC	I6L
P9	SAVYNFATC	C9A, C9D, C9E, C9F, C9K, C9L, C9M, C9N, C9R, C9S, C9V, C9W
P1, P9	SAVYNFATC	K1AC9M, K1MC9M, K1SC9M
P1, P9	SAVYNFATC	V1LC9M
P4, P9	SAVYNFATC	Y4FC9M, Y4SC9M

Table II. Effects of gp33 variants on T cell cytotoxicity, IFN-γ production, and TCR binding

Peptide	EC ₅₀ ^{CTL} (μM)	EC ₅₀ ^{IFN} (μM)	K _D (μM)	ΔG (kcal/mol)	k _{on} (×10 ⁶ M ⁻¹ s ⁻¹)	k _{off} (s ⁻¹)	k _{off} (s ⁻¹)
gp33	16.8 ± 16.8	0.96 ± 0.01	12.2 ± 2.2	-6.79	1.35 ± 0.27	1.35 ± 0.68	0.51
C9F	0.53 ± 0.36	0.56 ± 0.44	9.4 ± 0.5	-6.95	1.19 ± 0.01	1.15 ± 0.01	0.60
C9L	1.96 ± 2.1	0.47 ± 0.63	6.6 ± 0.2	-7.00	0.90 ± 0.02	0.79 ± 0.02	0.88
C9M	1.59 ± 2.67	0.92 ± 0.39	7.7 ± 1.7	-6.96	1.42 ± 0.54	1.2 ± 0.66	0.58
C9V	3.39 ± 5.00	1.23 ± 0.27	10.8 ± 0.3	-6.86	1.16 ± 0.03	1.25 ± 0.03	0.55
K1M/C9M	8.58 ± 5.27	0.13 ± 0.09	15.5 ± 3.1	-5.70	0.33 ± 0.08	0.53 ± 0.2	0.20
K1R	40.0 ± 22.6	7.08 ± 1.96	31.6 ± 10.9	-5.65	1.31 ± 0.47	10.6 ± 5.3	0.60
V1LC9M	>50,000	28.0 ± 4.55	207	-4.04	ND ^a	ND ^a	2.67
V1SC9M	>50,000	112.8 ± 19.3	261 ± 109	-4.64	ND ^a	ND ^a	ND ^a
Y4A	8.470 ± 5.550	11.27 ± 7.47	63.0 ± 14.3	-5.8	0.38 ± 0.07	2.2 ± 0.57	0.32
Y4SC9M	>50,000	38.83 ± 22.09	461	-4.58	ND ^a	ND ^a	ND ^a
Y4SC9M	>50,000	264.1 ± 157.5	550 ^b	-4.53	ND ^a	ND ^a	ND ^a

^aEC₅₀^{CTL} is the effective concentration of peptide that gives 50% maximum CTL activity as determined by 4 h chromium release assay. The values represent the mean ± SD of three to five experiments.

^bEC₅₀^{IFN} is the effective concentration of peptide that induces 50% of the maximum amount of IFN-γ as determined by ELISA. The values represent the mean ± SD of three to five experiments.

^cDerived from steady-state SPR data with software Scatchard. The values represent the mean ± SD of two to three experiments.

^dDerived from 30 s - 30 s K_D value from SPR data with software Scatchard.

^eQuantified from kinetic SPR data with software CLAMP 5.0. The values represent the mean ± SD of two to three experiments.

^fHalf-life of dissociation of pMHC from P14 TCR. Determined by SPR.

^gFrom kinetic data, k_{off} = k_{on} × K_D.

^hDerived from k_{off} = k_{on} × K_D.




FIGURE 3. Viral epitopes gp33 and APLs variably sensitize P14 T cells to kill target cells. ELA cells labeled with ⁵¹Cr and pulsed with peptides at various concentrations were incubated with activated splenocytes from P14 TCR transgenic mice (E:T ratio = 5) for 4 h. Specific lysis of ELA cells was measured by ⁵¹Cr release. Irrelevant [Y(K)A] peptide (ACSRNRYQL) was used as negative control. Data are shown as mean ± SD (n = 5), and are representative of three to five independent experiments. Using GraphPad Prism, data were fit into sigmoidal dose-response model to obtain EC₅₀^{CTL} values (shown in Table II) (fitting curves are not shown).

Importance de l'affinité du TCR pour l'activation lymphocytaire (2)





FIGURE 5. CTL response and IFN-γ production are correlated with binding affinity of peptide (D^b) to P14 TCR. a and b, Correlations between log EC₅₀^{CTL} and log K_D (a), log EC₅₀^{IFN} and log K_D (b). All the data points are shown in Table II. The EC₅₀^{CTL} and EC₅₀^{IFN} values are in unit of M, and K_D in μM. The solid lines are the fittings to Boltzmann sigmoidal equation as follows: Y = bottom + (top - bottom) / (1 + exp((V50 - X)/slope)), using GraphPad Prism. The K_D values for V1LC9M, Y4FC9M, and Y4SC9M are adopted from a previous report (42). Note that the EC₅₀^{CTL} values for K1SC9M, V1LC9M, Y4FC9M, and Y4SC9M were shown on the graph as 50 μM, although the actual values could be higher than that, but are out of the measurable range with CTL assay. c and d, Correlations between log EC₅₀^{CTL} and log K_D (c), log EC₅₀^{IFN} and log K_D (d). The solid lines in c and d are linear fittings using GraphPad Prism 4.01. All of the data points are shown in Table II. The EC₅₀^{CTL} and EC₅₀^{IFN} values are in unit of M.


L'affinité ne prédit pas toujours l'activation et la réponse lymphocytaire

- Activation:
 - Ont été observées :
 - Une diminution de l'activation liée à une augmentation de l'affinité
 - Une augmentation de l'activation liée à une diminution de l'affinité
- Spécificité

The Study of High-Affinity TCRs Reveals Duality in T Cell Recognition of Antigen: Specificity and Degeneracy
D. Donermeyer et al.; Journal of Immunology, 2006,



Degeneracy in the recognition of Hb correlates with increased affinity. CD4⁺ T cell hybridomas expressing 3.L2, M4, M14, and M15 TCRs were stimulated with the indicated peptide concentrations of Hb or APLs of Hb at the P8 (75) position using CH27 as APC. The APLs are designated as described in Fig. 2. The level of T cell stimulation was determined using a bioassay for IL-2. The cpm values of [3H]TdR incorporation into CTL-2 cells represent the mean ± SD of triplicate values of a representative experiment (n > 3).



Activation T : un système dynamique

Stimulation réitérative du lymphocyte: « kinetic proofreading » model

- Activation du lymphocyte T si interaction TCR/CMH-Ag suffisante (seuil d'activation)


Interactions :

- Force d'interaction (Affinité)
- Temps d'interaction (Koff)
- Nombre d'interactions (x n; Avidité)


- Activation progressive des molécules de signalisation (phosphorylation) : seuil d'activation

- Seuil critique pour initier une activation fonctionnelle


- Si interaction insuffisante, (dissociation du complexe TCR/CMH-Ag) avant initiation de l'activation: les groupements phosphates seront éliminés par les phosphatases cellulaires (McKeithan 1995)




Variabilité du niveau du seuil d'activation: T naïf vs T mémoire



Seuil d'activation: T naïf vs T mémoire




Beads H-2Kb+Ag




Proliferation of OT-1 CD8⁺ LNC in response to Kb/OVA₂₅₇₋₂₆₄ and rIL-2 is predominantly due to the CD44^{high} subset of cells. A, LNC (5 x 10⁴-sorted CD44^{high}CD8⁺ and CD44^{low}CD8⁺) from OT-1 mice were cultured with 1 x 10⁵ latex beads coated with H-2K^b at 1.7 μg protein/10⁷ beads and pulsed with OVA₂₅₇₋₂₆₄ peptide at the indicated concentration.

Dynamique de l'interaction Affinité versus Avidité



- Contacts multiples avec complexes CMH-Ag à la surface des DC



Correlation Between the Number of T Cell Receptors Required for T Cell Activation and TCR-Ligand Affinity




Figure 4. Relationship of Number of Ligands per Target Cell and the Number of TCRs Required for Activity. The number of 2C TCR required for recognition of various ligands was plotted as a function of the number of ligands on the target cell. The number of p2Ca-L¹, QL9-L¹, QL9-D5-L¹, SEB-class II, or SECI-class II complexes was calculated from the K_D of the peptide or ligand for MHC and the concentration of peptide or ligand added in the assay. The number of TCR required for recognition was determined from the concentration of 100-fold fragments required at 50% inhibition of cytokinesis, using a K_D of 2×10^5 M⁻¹ and 100,000 TCRs per T cell, as described in Experimental Procedures. Data are taken from Table 2.





Figure 5. Correlation Between the Binding Affinity of the TCR for its Ligand and the Number of TCRs Required for T Cell Activity. (A) K_D values from previous reports are summarized in Table 1. The number of TCR at >25,000 ligands (open circles) was determined at saturating ligand concentrations for each of the peptides or superantigens that have a published binding affinity for the 2C TCR. The number of TCRs at 1,000 (open triangles) and 10,000 (closed squares) ligands were calculated by regression analysis from the data summarized in Table 2. (B) K_D values for the TCR interaction with the indicated peptide-class II complexes were predicted from the number of TCRs required for activity. The curve determined from known K_D values for several TCR-ligand interactions was plotted versus number of TCRs at saturating ligand concentrations (open circles) to generate a standard curve. The inhibition analysis was used to determine the number of TCRs required at 5×10^{-5} M QL9-H5, QL9-R5, QL9-H5 and 10^{-5} M QL9-D5 with T2-L¹ and 10^{-5} M QL9 with T2-K¹. These values were plotted on the standard curve (closed circles) to calculate by linear regression a predicted K_D value for TCR binding to each of the peptide-class II complexes.



Modèle:
Clone 2C anti-AlloAg
Spé p2Ca/Ld; QL9/Ld
Nb ligands: Δ[peptide]
Nb TCR: [Fab 1B2],
bloque TCR
Affinité: Mutants de QL9

Nombre d'interactions TCR/CMH-Ag: aspects moléculaires

- 1 à 50 complexes CMH-Ag sur APC pour activer LyT
- Engagement multiple des TCR pour un complexe CMH-Ag
 - N= 200 contacts répétitifs (Valitutti 1995)
- Temps de contact cellulaire requis ? (Long / Court)
- Temps de contact TCR/CMH-Ag limité pour assurer des engagements répétés des TCR
- Engagement répété de la même molécule de TCR vs différentes molécules de TCR ?



Serial triggering of many T-cell receptors by a few peptide-MHC complexes Valitutti, Nature 1995




T cells conjugated with peptide-pulsed APCs undergo an antigen-dependent downregulation of the TCR/CD3 complex.

>> internalisation des complexes moléculaires « TCR »

Nombre d'engagements


Serial triggering of many T-cell receptors by a few peptide-MHC complexes
Valitutti, Nature 1995



we measured at different antigen concentrations the number of complexes per APC and the number of TCRs downregulated after T-APC interaction. In one series of experiments, EBV-B cells were pulsed with different concentrations of **125 I-labelled peptide** and the number of peptide-DR complexes per cell was calculated at each peptide concentration Figure 3(a). In parallel experiments, the fraction of TCRs downregulated by APCs pulsed in the same conditions was measured Figure 3(b). From comparison of the two curves it is estimated that **APCs pulsed with high peptide concentrations (20 micromolar) display approximately 7,500 complexes and induce downregulation of 93 percent of TCRs**. Strikingly, APCs pulsed with a low peptide concentration (50 nM) display only approximately 100 peptide-DR complexes, yet they downregulate 62 percent of TCRs. The relationship between the number of peptide-DR complexes per APC and the number of TCRs downregulated per T cell is shown in Figure 3(c). This plot clearly shows that each peptide-DR complex must engage a large number of TCRs in successive rounds. This effect is dramatic at low complex density, where approximately **100 complexes can trigger up to 18,000 TCRs**, but is less marked at high complex density, indicating that **a single peptide-DR must be able to trigger 180 TCRs in successive rounds**. This figure may increase at lower complex density and could be an underestimate as it is unlikely that all complexes present on an APC may be available to the responding T cell.

Internalisation du TCR

Coombs et al 2002



TCR down-regulation as a function of TCR-pMHC half-life.
Data are derived from 5-h TCR down-regulation experiments done with a panel of TCR mutants derived from a Kb-VSV-specific TCR (N30.7). Each point represents the indicated TCR-pMHC pair: 1, V98D-Kb-VSV; 2, V98L-Kb-VSV; 3, N30.7-KbA158T-VSV; 4, G99A-KbA158T-VSV; 5, G97/99A-KbA158T-VSV; 6, G97A-KbA158T-VSV; 7, N30.7-Kb-VSV; 8, G97A-Kb-VSV; 9, G97/99A-Kb-VSV; 10, G99A-Kb-VSV. Half-lives are shown as relative units (RU) normalized to the half-life for the N30.7-Kb-VSV (number 7) interaction. TCR down-regulation data are shown as RU normalized to TCR down-regulation measured for N30.7 hybridomas that were interacting with 10⁻⁴ - 10⁻⁵ M VSV peptide.

TCR internalization as a function of the dissociation rate constant.
Results are shown for the monomer model (closed circles) and the dimer model (open squares). (a) Only activated bound TCRs are internalized, lambda_B = lambda_D = 0.003 s⁻¹, lambda_T = 0 s⁻¹. (b) Only unbound activated TCRs are internalized, lambda_B = 0 s⁻¹, lambda_T = 0.003 s⁻¹.

> Internalisation du TCR dépend:
-De la constante de dissociation
-De temps de contact

> Après libération du CMH-Ag = Modèle 2

Interactions en série





Fig. 3. Parameters that influence TCR triggering and signaling at the level of the T-cell-APC interaction. Sustaining the process requires a continuous supply of signal transduction components from the cytosol, as well as new TCRs to substitute for those that are downregulated and degraded¹⁰. Cell-cell adhesion is boosted by TCR signaling and forces TCR engagements with peptide-MHC complexes. Abbreviations: see Fig. 1 legend.

Serial triggering of TCRs: a basis for the sensitivity and specificity of antigen recognition

Salvatore Valitutti and Antonio Lanzavecchia

Serial triggering of TCRs: a basis for the sensitivity and specificity of antigen recognition

Salvatore Valitutti and Antonio Lanzavecchia





Fig. 2. Serial TCR triggering or inactivation as a function of complex stability and time required for TCR triggering. (a) Simulation describing the number of TCRs triggered in 30 min as a function of the half-life of TCR-peptide-MHC interactions. Each curve is calculated for times required for TCR triggering ranging from 1 to 15 s. (b) Simulation describing the number of TCRs fully triggered (green), partially triggered (blue) or inactivated (red) as a function of the half-life of TCR-peptide-MHC interaction. It is assumed that ligand for 20 s results in full TCR triggering, ligand for 5-15 s in partial triggering and ligand for 1-3 s in TCR inactivation. Abbreviations: see Fig. 1 legend.


	Agonist	Antagonist
T1/2	3-10	0.5-1
#TCR	100	250

Paramètres d'activation



Importance de la synapse immunologique pour favoriser la multiplication des contacts

- Favorise interactions multiples TCR/CMH-Ag
- Rôle dans la transduction du signal ?
 - Formation de la synapse est associée à l'activité PI3K
 - Blocage de la synapse empêche signal calcique (tardif) et la production de cytokine (IFN γ) mais ne perturbe pas les événements de signalisation précoces (phosphorylation ERK)
- Lien étroit entre activation T et maintien de la structure « synapse »
 - De la même façon qu'il existe des paramètres d'interaction du TCR optimum, la formation des suprastructures cSMAC/pSMAC dépend de la demi-vie d'interaction du TCR



Recruitment and activation of proximal signalling intermediates — such as lymphocyte-specific protein tyrosine kinase (LCK), ZAP70 (chain-associated protein of 70 kDa), LAT (linker for activation of T cells), SLP76 (SRC homology-2-domain containing leukocyte protein of 76 kDa), NCK (non-catalytic region of tyrosine kinase), ITK (interleukin-2-inducible T-cell kinase) and VAV1—is probably initiated at sites of T-cell receptor (TCR)-microcluster formation^{80, 84}. Recruitment of these molecules leads to the activation of filamentous (F)-actin-regulatory proteins, which control actin polarization in T cells. Interestingly, signals that arise from TCR microclusters not only induce F-actin polymerization, but are required for the stable formation and movement of TCR microclusters, and ultimately lead to the organization of the mature immunological synapse (consisting of the central and peripheral supramolecular activating clusters; cSMAC and pSMAC, respectively)^{13, 29, 80}. In addition, global F-actin reorganization resulting from T-cell activation leads to lamellipodium formation and distal-pole formation, facilitating T-cell-APC (antigen-presenting cell) recognition¹¹. DLG1 (disc-large homologue-1)¹³, ERM (ezrin, radixin and moesin) and F-actin also polarize to the rear of the T cell to form the distal-pole complex^{11, 48}. It is thought that this complex recruits some receptors to the rear of the cell, away from the immunological synapse.

Importance de la « demie-vie » de contact




Fig. 2. T cell activation depends on TCR/pMHC interaction half-life and pMHC density. (A) When T cells interact with cognate pMHC at low density on the APC, efficient activation takes place within an optimal range of TCR/pMHC interaction half-life. Interactions with short half-lives cannot complete necessary intracellular signals for T cell activation, due to impairment on TCR kinetic proofreading. TCR/pMHC interactions with excessively long half-lives impair T cell activation due to TCR serial engagement blockade. At low pMHC density, the plot of T cell activation versus TCR/pMHC half-life results in a Gaussian distribution, in which only pMHCs that interact with intermediate half-lives with the TCR behave as agonists (Kaleris et al., 2001; Coombs et al., 2002). (B) When T cells interact with cognate pMHC at high density on the APC, T cell activation can take place when the half-life of the TCR/pMHC interaction is intermediate or high, because serial engagement requirement no longer applies (Gonzalez et al., 2005). At high pMHC density, the plot of T cell activation versus TCR/pMHC half-life results in a sigmoid distribution, in which pMHCs that interact with intermediate and long half-lives with the TCR behave as agonists.

Conséquences sur l'immunodominance






Fig. 3. Clonal expansion of T cells in response to antigenic challenge is determined by the half-life of TCR/pMHC interaction. Before antigen challenge, the T cell population shows a wide range of half-lives for the TCR/pMHC interaction. After the first antigen challenge, only T cells bearing TCRs with optimal half-lives are clonally expanded and the global TCR/pMHC half-life of the responding T cells shifts towards an optimal value. Although subsequent antigenic challenges do not lead to a significant increase in the overall TCR affinity of the T cell population, there is a selection of T cells within a narrower range of TCR/pMHC half-lives.




la diminution d'expression des CMH limite la réactivité des T
Cas de l'échappement viral et tumoral:

A review of viral strategies to evade selected components of the host immune system, Bruce Wobeser




Virus	Gene / Gene product	Function
Adenovirus	E1A ₁	Inhibition of MHC-I in ER
Adenovirus	Adenovirus hexon	Disruption of trans-Golgi network
Bovine Papilloma Virus	E5	Retention of MHC-I to plasma membrane
Coxsack Virus	Unknown early gene product	Inhibition of MHC-I expression
Human Cytomegalovirus	Unknown early gene product	HLA-DR segregation and phosphorylation of T2-44 translocator
	US2	Disruption of MHC-I heavy chain
	US3	Retention of MHC-I to TAP
	US6	Inhibition of TAP
	US11	Disruption of MHC-I and CD4
	US4	Retention of MHC-I in TAP
	US7	Inhibition of MHC-I and CD4
	US9	Inhibition of MHC-I
	US10	Disruption of newly synthesized MHC-I
	US12	Inhibition of TAP
Human Papilloma Virus	Unknown	Inhibition of TAP
Human Papilloma Virus	Unknown	Inhibition of MHC-I gene promoter, retention of TAP gene promoter
Herpes Simplex Virus	ICP-47	Inhibition of TAP gene promoter
Kaposi's Sarcoma Herpes Virus	ICP-47	Targets MHC-I for degradation
Marburg Cytomegal Virus	K5	Retention of MHC-I in ER / Golgi
Measles Cytomegal Virus	NS2	Retention of MHC-I in ER / Golgi
Murine Cytomegal Virus	pp65	Targets MHC-I for degradation
Respiratory Virus	NS1	Retention of MHC-I in ER / Golgi
Rubella Virus	n133R	MHC-I internalization and degradation
Schistosoma	Unknown	Disrupts signal pathway of MHC-I expression
Symptomatic Virus	Unknown	Inhibition of MHC-I expression



Tumor escape from immune recognition: loss of HLA-A2 melanoma cell surface expression is associated with a complex rearrangement of the short arm of chromosome 6 MJ Maeurer, MJ Maeurer et al; Clinical Cancer Research, 1996

Importance des paramètres de fixation du TCR en fonction de la fréquence des complexes CMH-Ag



Sélection clonale:
immunodominance


Importance de la sur-expression des molécules de CMH par les dendritiques après activation

The temporal TCR signal summation model suggests the existence of an optimal TCR/pMHC half-life for T cell activation in response to low-density cognate pMHC ligand. (A) TCR/pMHC interactions with short half-lives fail to activate the T cell (inset) in response to low-density pMHC due to summation of TCR signalling not reaching the activating threshold. (B) TCR/pMHC interactions with optimal half-life lead to efficient T cell activation (inset) as a result of summation of productive TCR signalling that reaches the activating threshold. (C) TCR/pMHC interactions with prolonged half-lives fail to activate T cells (inset) in response to low-density pMHC, due to TCR signals being generated at low frequency (TCR serial engagement blockade) and fail to be added one over the other due to intrinsic signal decay. Thus, T cell activation is impaired (inset) as activating threshold is not reached. (D) However, TCR/pMHC interactions with prolonged half-life can lead to T cell activation (inset) in response to high-density pMHC, because multiple productive TCR signals are generated simultaneously. These signals can be added and the T cell activation threshold reached.

Paramètres quantitatifs de l'activation

10⁵ TCR / T

**Engagement de
à 400 TCR différents**



x 1- 200 / t = 2 - 30 s



Dépend de la présence de

CD8

CD28
CD40L


LFA-1

- **Molécules adaptatrices:**
 - Stabilisation de l'interaction TCR/CMH
 - Modèles d'initiation de l'activation
- **Molécules de costimulation:**
 - Transduction du signal : **renforcement des voies de signalisation**






Michelle Krogsqaard, Sim Imm 2007

Kerry JI 2003




- **Model:**
- TCR-transgenic CD8 (P14)
- GP33 (KAVYNFATC) ou C9M (KAVYNFATM; affinity >) peptides presented by
- H-2Db tetramers or tetramers containing the **D^b D227K** mutation, which has been shown to **abrogate CD8** interaction with MHC




Holler D , Imm 2003

Transfection TCR +/- CD8



The significant enhancement in recognition of the peptides could be improved by one to two orders of magnitude by the introduction of CD8 into transfectants




The $\alpha\beta$ -negative T cell hybridoma 58-/- was cotransfected with the 2C wt α chain gene and one of five different α chain genes from 2C and the mutants m6 α m13 α m33 α and m67 α . Stable transfectants that expressed similar levels of TCR were identified based on their staining with anti-V α antibody KJ16

Sensitization Doses of Various QL9 Position 5 Variant PeptidesIL-2 production by transfectants stimulated with various peptides was measured as described in the Experimental Procedures. The amount of peptide that yielded 50% of the maximum IL-2 release (SD50) was calculated by linear regression of IL-2 curves (see Figure S1

at <http://www.immunity.com.gate1.inist.fr/cgi/content/full/18/2/255/DC1>). The log of the SD50 value was plotted for each of the peptides used to stimulate transfectants: (A) 2C and m6₈-CD8-negative; (B) 2C-CD8-negative, 2C/CD8₈ and 2C/CD8₈-(C) m6₈-CD8-negative, m6₈-CD8₈ and m6₈-CD8₈-. In [Figure 3](#), [Figure 4](#) and [Figure 5](#), log SD50 values represent the mean of at least two assays. Error bars indicating one standard deviation are included for each point (for some points these error bars are not apparent as they are smaller than the size of the symbol).

Relationship of Peptide Activity and TCR-peMHC Affinity. S050 values of various peptides were plotted versus the equilibrium affinity constants (KD) of the corresponding TCR-peMHC interaction (Table 1). In order to include values from the Ld and Kb systems, S050 values from the corresponding TCR-pMHC interaction were also included (Table 2-Kb) are shown on the two Y-axes. The points represent data derived from CD8-negative transfectants (blue circles) and CD8+ transfectants (red squares) of the corresponding TCR transfectant. The corresponding TCR-pMHC interaction is indicated by the dashed line in the figure. The range of KD values that correspond to affinities measured for known TCR-syngeneic MHC interactions from *vivo* CTLs are shown. This range falls exclusively in the CD8-dependent category, as determined by TCR transfection studies (not necessarily anti-CD8 antibody inhibition studies).


- **Absence:**
 - mécanisme de tolérance périphérique: ignorance immunologique
 - Induction d'anergie



Mécanismes possibles de la tolérance périphérique


Figure N°22

- Mol de costimulation: signal complémentaire



Nécessité d'un engagement prolongé de CD80/CD86

Liwski, Imm Letters 2006




- To investigate the functional effects of CD80/CD86 blockade on naive CD4⁺ T cell activation, freshly isolated CD4⁺ T cells from OVA-TCR transgenic DO11.10 mice [27] were labeled with CFSE and cultured with OVA-peptide pulsed, mature DC in the absence or presence of CD80 and CD86 specific mAbs.



- We demonstrated that DC mediated CD80/CD86 costimulation controls the magnitude of the naive T cell proliferative response by regulating both responder frequency as well as proliferative capacity.
- Our data revealed that blocking CD80/CD86 signaling up to 6 h after conjugate formation resulted in a significant decrease in both the number of naive T cells entering the proliferative cycle and the number of daughter cells generated by each cell.

Mol costimulation et dynamique de l'interaction

- Federica Benvenuti, JI 2004; Dendritic Cell Maturation Controls Adhesion, Synapse Formation, and the Duration of the Interactions with Naive T Lymphocytes




- Proliferation of naive T cells stimulated by immature or mature DCs. Immature or mature D1 pulsed with different doses of peptide were cocultured for 5 days with CFSE-loaded naive T lymphocytes (1:5 ratio). A, Representative dot blot profile showing the loss of CFSE and the up-regulation of CD44 induced by immature (upper row) or mature (lower row) DCs loaded with different peptide doses at day 3. B, Histogram profile of CFSE staining on naive T cells stimulated with immature (upper row) or mature (middle row) DCs loaded with 1 nM H-Y and mature DCs loaded with 0.1 nM peptide (lower row) at days 2-5 of the coculture. C, Quantification of the absolute number of CD4⁺ T cells at the different days of coculture for immature (○) and mature (●) DCs loaded with 1 nM peptide (T cells at day 0 = 7 × 10⁴). One representative of three experiments is shown.



Mol de costimulation et synapse

- Federica Benvenuti, JI 2004; Dendritic Cell Maturation Controls Adhesion, Synapse Formation, and the Duration of the Interactions with Naive T Lymphocytes



Conséquence des interactions T sur la différenciation T

DC maturation is required for effective clustering and SMAC formation at the DC-T cell contact site

Efficient clustering in naive T cells requires DC maturation. Confocal images showing the distribution of CD3, LFA-1, LAT, and tubulin in T cells forming conjugates with immature or mature D1 pulsed with 10 nM H-Y peptide. Conjugates were formed for 30 min, washed five times, and fixed for immunostaining. A, One representative conjugate formed with immature DCs (upper panels) or mature DCs (lower panels) is shown for each marker. For each immunofluorescent image (right panels), a DIC image showing the two cells in contact is shown (left panels). Note that the distribution of CD3, LFA-1, and LAT is homogenous on T cells forming conjugates with immature DCs and clustered in conjugates formed with mature DCs. Similarly, the T cell MTOC (marked by an arrow) is reoriented toward the APCs in conjugates with mature, but not with immature DCs. B, Quantification of the proportion of conjugates showing clustering of CD3, LFA-1, LAT, and tubulin at the site of contact. Conjugates between naive T cells and immature or mature DCs pulsed with different doses of H-Y peptide were formed as in A. The number of conjugates presenting clustering at the contact zone was blindly quantified (see Materials and Methods) and expressed as a percentage of the total number of conjugates analyzed (percent clustering). Number of conjugates quantified is as follows: CD3, immature, n = 276; mature, n = 347; LFA-1, immature, n = 197; mature, n = 229; LAT, immature, n = 185; mature, n = 232; and tubulin, immature, n = 210; mature, n = 243.

Génération des CD8 mémoire: Influence de la quantité d'Ag


- E. John Wherry, *Jl* 2002, Generation of CD8+ T Cell Memory in Response to Low, High, and Excessive Levels of Epitope

Table I. VV recombinants


Recombinant	Description	Epitopes/Cell* (copies/cell)
M50	Control virus-expressing an irrelevant minigene (NP ₂₂₂₋₂₂₇ : H-2K ^b)	0
HP19	Full-length NP/S (influenza NP + SIINFEKL) behind a 19bp HP	~800
HP0	Full-length NP/S (influenza NP + SIINFEKL) without a HP (HP size = 0)	~30,000
(M)SIINFEKL	Minigene version of OVA ₂₃₇₋₂₄₄ (M designates the initiating methionine)	~60,000

Espaceur:
< expression
de l'Ag

C Kb/SIINFEKL-Specific Cells/Spleen by MHC/Tetramer Staining



A SIINFEKL-Specific IFN-γ Secreting Cells/Spleen by ELISPOT




➤recrutement d'un nombre de clones plus ou moins important


➤et d'affinité variable

selon quantité d'Ag lors de la stimulation

Prolongation de la stimulation antigénique: conséquence sur la différenciation en cellules mémoire



Importance des paramètres d'activation : Génération des CD8 mémoire





Masopust Curr Op Imm 2004

The Goldilocks model of effector and memory CD8+ T-cell development. The first 24 hours following antigenic stimulation elicits a program of expansion and differentiation that continues among daughter cells after removal of antigen. 'Too little' stimulation, meaning insufficient antigen concentration or duration, leads to limited CD8+ T-cell expansion, poor memory development and attrition. Chronic antigen exposure may cause 'too much' stimulation, leading to a progressive loss in the ability to secrete cytokines and the eventual deletion of antigen-specific CD8+ T cells. Optimum memory development is favored when conditions are 'just right'; that is, when CD8+ T cells are stimulated by a sufficient concentration of antigen for a sufficient, but not excessive period of time. Several variables, such as the cytokine environment, TCR affinity, type of antigen presenting cells, CD4+ T-cell help and degree of co-stimulation, modify the differentiation program and alter the antigen-dependent signaling requirements for the development of a robust response. Several factors encountered during the days following priming, including continued antigen contact, various co-stimulatory molecules, negative regulators, cytokines, chemokines, CD4+ T-cell help and anatomical location might also have a large qualitative and quantitative influence on the developing response. Additional factors may be important weeks and months later for memory differentiation and maintenance.

Importance de la stimulation antigénique initiale: programme de différenciation

- Susan M. Kaech & Rafi Ahmed *Nature Immunology* 2001 ; Memory CD8+ T cell differentiation: initial antigen encounter triggers a developmental program in naive cells
- Models for proliferation and differentiation of naive CD8+ T cells. (a) CD8+ T cell proliferation is dependent on repeated encounters with antigen. Each cell that is stimulated by antigen divides and progressively differentiates into effector CTLs and memory CD8+ T cells with each successive cell division. According to this model, it is essential that each daughter cell be stimulated with antigen, otherwise CD8+ T cell division, and possibly differentiation, would be halted upon antigen removal. (b) Naive CD8+ T cells are developmentally programmed to divide at least seven to ten times and to differentiate into effector CTLs and long-lived functional memory CD8+ T cells. Optimal antigenic stimulation of the parental cell triggers this developmental program and the CD8+ T cells become committed to proliferation and differentiation. Further antigenic stimulation of the daughter cells may increase the number of times the activated CD8+ T cells divide, but it is unnecessary for this developmental program to progress.





Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells

Kaeche, Nat Immunol 2003

IL-7R^{hi} memory cell precursors provide immunological protection

Thy-1.1⁺ P14 effector cells (2 × 10⁵) from day 11 were adoptively transferred into naive hosts, and 3 weeks later the mice were infected with 5 × 10⁶ CFU of recombinant *L. monocytogenes* expressing the GP33-41 epitope (LM-GP33). Mice were examined 5 d later. (b) Donor Thy-1.1⁺ P14 cell population expansion. Dot plots gated on CD8 T cells; percent indicate Thy-1.1⁺ P14 cells in the spleen. (c) Bacterial clearance. Bacterial titers were measured in spleen and liver. Similar results were obtained in separate experiments in which IL-7R^{hi} and IL-7R^{lo} effector cells from day 15 were transferred. (d) Equal numbers (0.8 × 10⁶) of IL-7R^{hi} or IL-7R^{lo} effector Thy-1.1⁺ P14 CD8 T cells from day 8 after infection (d8) or memory Thy-1.1⁺ P14 CD8 T cells from day 40 or later after infection (d40+) were transferred into mice persistently infected with LCMV, and their ability to proliferate *in vivo* after antigenic stimulation was assessed. Data represent percentage of donor Thy-1.1⁺ P14 cells of total CD8 T cells in the blood on days 5, 8 and 15 after transfer.

Increased expression of IL-7R directly correlates with effector CD8 T cell survival


(a) Differential expression of IL-7R mRNA in naive, effector and memory P14 CD8 T cells. Expression of IL-7R mRNA was measured by DNA microarray analysis in P14 CD8 T cells from days 8, 15, 22 and >40 after LCMV infection and was compared with that of naive P14 CD8 T cells. Bars represent the relative fold difference based on a log scale. The P14 CD8 T cells express a transgenic TCR that recognizes the GP33-41 epitope of LCMV. (b) Cell surface IL-7R expression on naive P14 CD8 T cells or P14 CD8 T cells from days 8, 15, 22 and >40 after infection. The percent and mean fluorescent intensity (MFI) of bracketed IL-7R^{hi} cells are given. (c) C57BL/6 mice were infected with LCMV, and 8, 15, 22 and >40 d after infection, the cell surface expression of IL-7R on pGP33-41- and pNP396-404-specific CD8 T cells was assessed by staining with anti-CD8 and anti-IL-7R and either pGP33-41 or pNP396-404 major histocompatibility complex class I tetramers. The dot plots are gated on tetramer-positive CD8 T cells, and the percent of IL-7R^{hi} cells is indicated. Circles and arrows indicate the source of cells in dot plots to the right. (d) LCMV-specific CD8 T cells in the spleen 8, 15, 22 and >40 days after infection were quantified by major histocompatibility complex class I tetramer staining as well as by intracellular cytokine staining for IFN-γ after 5 h of stimulation with LCMV peptides. The percent of LCMV-specific CD8 T cells predicted to survive at day 8, 15, 22 and approximately 40 days after infection was calculated by dividing the number of LCMV-specific memory CD8 T cells (at day 40 or more after infection) by the number of virus-specific CD8 T cells at each time point. (e) C57BL/6 (Thy-1.2⁺) mice containing 1 × 10⁶ naive TCR-transgenic CD8 T cells from Thy-1.1⁺ P14 mice that recognize the LCMV epitope GP33-41 were infected with LCMV, and 8 d later the expression of granzyme B, CD62L, CD43, CD27 and KLRG-1 was compared between IL-7R^{hi} (blue) and IL-7R^{lo} (red) Thy-1.1⁺ effector cells and naive CD8 T cells (black). Arrow indicates the source of cells in histograms.

UPMC PARIS UNIVERSITÉS B. Bellier

Effector CD8 T cells uniformly down-regulate CD127 but can be distinguished on the basis of KLRG-1 expression. (A) Cell-surface expression of IL-7R with respect to cell division at days 104 after LCMV infection. 10⁶ naive CFSE-labeled P14 CD8 T cells were adoptively transferred into naive mice that were subsequently infected with LCMV. All plots are gated on CD8⁺ Thy1.1⁺ P14 splenocytes directly stained ex vivo for IL-7R expression. An uninfected naive control is also shown, and the red horizontal line indicates the naive level of expression. (B) Phenotypic properties of CD127^{hi} antigen-specific cells at day 4.5 p.i. B6 mice containing ~10⁵ naive Thy1.1⁺ P14 cells were infected with LCMV, and expression of the indicated cell-surface and intracellular markers on CD8⁺ Thy1.1⁺ splenocytes was assessed 4.5 d later (red line histograms). Gray histograms represent naive cells from uninfected control mice. (C) Longitudinal analysis of cell-surface IL-7R and KLRG-1 expression. B6 mice containing ~10⁵ naive Thy1.1⁺ P14 cells were infected with LCMV, and the cell-surface expression of IL-7R and KLRG-1 was analyzed at the indicated time points. Histograms depict the MFI of IL-7R expression and percentages of KLRG-1^{hi} cells gated on CD8⁺ Thy1.1⁺ P14 cells. (D) Inverse correlation of KLRG-1 expression with the memory markers CD127 and CD62L. B6 mice containing 10⁶ CD8⁺ Thy1.1⁺ P14 cells were infected with LCMV, and KLRG-1 expression with respect to CD127 or CD62L was analyzed at the indicated time points.


Functional and genomic profiling of effector CD8 T cell subsets with distinct memory fates

Sarkar S et al, JExpMed 2008




SLEC KLRG1^{hi} IL-7R^{lo} MPEC KLRG1^{hi} IL-7R^{hi}

Short effector / Memory precursor modulation by T-bet




Inflammation Directs Memory Precursor and Short-Lived Effector CD8(+) T Cell Fates via the Graded Expression of T-bet Transcription Factor
Joshi et al, Immunity 2007




Levels of inflammation regulate KLRG1^{hi} IL-7R^{hi} SLEC formation.

UPMC PARIS UNIVERSITÉS B. Bellier

Model of SLEC and MPEC development during acute viral infection.



Naive CD8 T cells are IL-7R^{hi}, CD122^{hi} (IL-2/15^{hi}), KLRG1^{hi} and T-bet^{hi} and are IL-7 dependent. Early during infection, most effector CD8 T cells become CD122^{hi} and downregulate IL-7R to an intermediate-to-low level, but expression of T-bet and KLRG1 is set depending on their exposure to inflammatory cytokines (e.g. IL-12). Effector CD8 T cells that are exposed to lower levels of inflammation express less T-bet (light blue cells) and begin to upregulate IL-7R to become KLRG1^{hi} IL-7R^{hi} MPECs (turquoise cells). Effector CD8 T cells that encounter higher levels of inflammatory cytokines express relatively more T-bet and KLRG1 (dark blue cells), stably repress IL-7R and consequently become KLRG1^{hi} IL-7R^{lo} SLECs. SLECs become IL-15 dependent, however, IL-15 alone cannot support their long-term persistence or homeostatic turnover and they decline over time. In contrast, MPECs remain dually responsive to IL-7 and IL-15 and preferentially develop into long-lived memory CD8 T cells that can self-renew.



Tbeth^{hi}



Prot (degrad Tbet)

Eff

Mem

unpublished

Différenciation dès la 1ere division



- Paramètres de l'interaction TCR/CMH-Ag
 - Affinité
 - Avidité
 - Temps contact
 - Flexibilité du CMH-Ag
 - Molécules adaptatrices / costimulation

- Déterminent
 - Activation lymphocytaire
 - Différenciation lymphocytaire

Conclusions

